

ACETAMINOPHEN TOXICITY

BASICS

OVERVIEW

- Results from owners overdosing the patient with over-the-counter medications containing acetaminophen, a medication intended to control pain or fever in humans

GENETICS

- Cats—genetic deficiency in a pathway that breaks down or changes (metabolizes) drugs in the liver (known as the “glucuronide conjugation pathway”); makes cats vulnerable to acetaminophen toxicity

SIGNALMENT/DESCRIPTION of ANIMAL

- Cats; dogs
- Most common drug toxicity in cats; considerably less frequent in dogs
- Young and small dogs and cats—greater risk from owner-given single-dose acetaminophen medications

SIGNS/OBSERVED CHANGES in the ANIMAL

- May develop 1–4 hours after dosing
- Progressive depression
- Rapid breathing
- Darkened mucous membranes (moist tissues of body, such as gums)
- Drooling (salivation)
- Vomiting
- Abdominal pain
- Rapid breathing (tachypnea) and bluish discoloration of skin and moist tissues of body (cyanosis) due to a abnormal compound (methemoglobin) in the blood (methemoglobinemia) that disrupts the ability of the red blood cells to carry oxygen to the body
- Fluid build up (edema)—face, paws, and possibly forelimbs; after several hours
- Chocolate-colored urine due to the presence of blood in the urine (hematuria) and the presence of methemoglobin in the urine (methemoglobinuria); especially in cats
- Death

CAUSES

- [Acetaminophen](#) overdosing

RISK FACTORS

- Nutritional deficiencies of glucose and/or sulfate
- Simultaneous administration of other glutathione-depressing drugs

TREATMENT

HEALTH CARE

- With methemoglobinemia—must evaluate promptly; inpatient care
- With dark or bloody urine or yellowish discoloration of skin and moist tissues of the body (jaundice or icterus)—inpatient care
- Gentle handling—imperative for clinically affected patients
- The veterinarian will induce vomiting (emesis) and may perform flushing of the stomach (gastric lavage)—useful within 4–6 hours of ingestion of acetaminophen
- Low blood count (anemia), blood in the urine (hematuria), or presence of hemoglobin in the urine (hemoglobinuria)—may require whole blood transfusion
- Fluid therapy to maintain hydration and electrolyte balance
- Drinking water should be available at all times

ACTIVITY

- Restricted

DIET

- Food—offered 24 hours after initiation of treatment

MEDICATIONS

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all inclusive.

- **Activated charcoal**—administered immediately after the veterinarian has induced vomiting or flushed the stomach (gastric lavage) and after vomiting is controlled; activated charcoal is used to attract and keep the remaining acetaminophen in the gastrointestinal tract
- *N*-acetylcysteine (Mucomyst) is administered; considered to be an antidote for acetaminophen toxicity
- Other sulfur donor drugs—if *N*-acetylcysteine not available; sodium sulfate
- 1% methylene blue solution—combats methemoglobinemia without inducing red blood cell destruction (known as a “hemolytic crisis”)
- Ascorbic acid—slowly reduces methemoglobinemia

FOLLOW-UP CARE

PATIENT MONITORING

- Continual clinical monitoring of methemoglobinemia
- Serum liver enzyme activities to monitor liver damage
- Blood glutathione level—provide evidence of the effectiveness of therapy

PREVENTIONS AND AVOIDANCE

- Never give acetaminophen to cats
- Give careful attention to acetaminophen dose in dogs; acetaminophen should only be given to dogs under a veterinarian’s supervision

POSSIBLE COMPLICATIONS

- Liver damage and resulting scarring (fibrosis)—may compromise long-term liver function in recovered patients

EXPECTED COURSE AND PROGNOSIS

- Rapidly progressive methemoglobinemia—serious sign
- Methemoglobin concentrations greater than 50%—grave prognosis
- Progressively rising serum liver enzymes 12–24 hours after ingestion—serious concern
- Expect clinical signs to persist 12–48 hours; death owing to methemoglobinemia possible at any time
- Dogs and cats receiving prompt treatment that reverses methemoglobinemia and prevents excessive liver damage may recover fully
- Dogs—death as a result of liver damage may occur in a few days
- Cats—death as a result of methemoglobinemia occurs 18–36 hours after ingestion

KEY POINTS

- Never give acetaminophen to cats
- Acetaminophen should only be given to dogs under a veterinarian’s supervision
- Most common drug toxicity in cats; considerably less frequent in dogs
- Treatment in clinically affected patients may be prolonged and expensive
- Patients with liver injury may require prolonged and costly management

